

WEIGHTED DISPERSIVE ESTIMATES FOR TWO-DIMENSIONAL SCHRÖDINGER OPERATORS WITH AHARONOV-BOHM MAGNETIC FIELD

GABRIELE GRILLO AND HYNEK KOVAŘÍK

ABSTRACT. We consider two-dimensional Schrödinger operators H with Aharonov-Bohm magnetic field and an additional electric potential. We obtain an explicit leading term of the asymptotic expansion of the unitary group e^{-itH} for $t \rightarrow \infty$ in weighted L^2 spaces. In particular, we show that the magnetic field improves the decay of e^{-itH} with respect to the unitary group generated by non-magnetic Schrödinger operators, and that the decay rate in time is determined by the magnetic flux.

1. Introduction

Long time behavior of propagators of Schrödinger operators is known to be closely related to the spectral properties of their generator near the threshold of the continuous spectrum. For example, if zero is a regular point of a Schrödinger operator $-\Delta + V$ in $L^2(\mathbb{R}^n)$ in the sense of [JK], then in a suitable operator topology

$$e^{-it(-\Delta+V)}P_{ac} = \mathcal{O}(t^{-n/2}) \quad t \rightarrow \infty, \quad (1.1)$$

provided the electric potential $V : \mathbb{R}^n \rightarrow \mathbb{R}$ decays fast enough. Here P_{ac} denotes the projection onto the absolutely continuous spectral subspace of $-\Delta + V$. For $n = 3$ such dispersive estimates were established in [Ra, Je, JK] in weighted L^2 spaces. For corresponding $L^1 \rightarrow L^\infty$ bounds we refer to [JSS, GS, Wed] in the case $n = 1, 3$, and to [Sch1] in the case $n = 2$. Situations in which zero is not a regular point of $-\Delta + V$ are studied in great generality in [JK], see also [ES1, ES2].

On the other hand, very little is known about estimates of type (1.1) for magnetic Schrödinger operators, where $-\Delta + V$ is replaced by $(i\nabla + A)^2 + V$ with a vector potential A . Recently it was shown in [KK] that for $n = 3$ equation (1.1) can be extended to magnetic Schrödinger operators, in suitable weighted L^2 spaces, under certain decay and regularity conditions on A and V . As for the case $n = 2$, it was proved, see [FFFP], that (1.1) holds true when $V = 0$ and $A = A_{ab}$ is the vector potential generating the so-called Aharonov-Bohm magnetic field, see equation (2.1) below.

The aim of this paper is to point out the *diamagnetic effect* on the dispersive estimates (1.1) in the case $n = 2$. More precisely, we want to show that the magnetic field improves the decay rate of the propagator in dimension two. This is partially motivated by the following fact [Mu, Sch2, Go, EG]: if a Schrödinger operator $-\Delta + V$ in $L^2(\mathbb{R}^n)$ with $n = 1, 2$ does not have a resonance at zero energy, then the time decay of $e^{-it(-\Delta+V)}$, considered in suitable topology, is faster than the one predicted by (1.1). Since a magnetic field in \mathbb{R}^2 generically removes the resonance at zero energy, see [LW, W], it is natural to expect a *faster* time decay for propagators generated by magnetic Schrödinger operators with respect to the non-magnetic ones.

We will prove this conjecture in the case of the Aharonov-Bohm magnetic field, that is for operators of the type $(i\nabla + A_{ab})^2 + V$. In particular, we will show, under suitable assumptions on V , that in certain weighted L^2 spaces the associated propagator decays as

$$t^{-1-\min_{k \in \mathbb{Z}} |k-\alpha|}, \quad t \rightarrow \infty, \quad (1.2)$$

where α is the total flux of the Aharonov-Bohm field, see Theorem 3.6 for details. This is to be compared with the t^{-1} decay rate of non-magnetic Schrödinger propagators, see (1.1). In the special case $V = 0$ we obtain a stronger result which provides a point wise upper bound on the integral kernel of $e^{-it(i\nabla + A_{ab})^2}$, see Theorem 3.1 and Corollary 3.3.

Note that the decay rate of the unitary group generated by $(i\nabla + A_{ab})^2 + V$ is proportional to the distance between the magnetic flux and the set of integers, see (1.2). This is expected since an Aharonov-Bohm field with an integer flux can be gauged away and therefore does not affect the spectral properties of the corresponding generator. Our main results, Theorem 3.1 and Theorem 3.6 are presented in section 3. The proofs of the main results are given in section 6.

Although our paper deals only with the case of the Aharonov-Bohm field, we believe that a similar improved time decay should occur for a much wider class of magnetic fields, see Remark 3.9 for further discussion.

2. Preliminaries

The vector potential

$$A_{ab}(x) = (A_1(x), A_2(x)) = \frac{\alpha}{|x|^2} (-x_2, x_1) \quad \text{on} \quad \mathbb{R}^2 \setminus \{0\}, \quad (2.1)$$

generates the Aharonov-Bohm magnetic field which is fully characterized by its constant flux α . It is well-known, see e.g. [AT], that the operator

$$(i\nabla + A_{ab})^2 \quad \text{on} \quad C_0^\infty(\mathbb{R}^2 \setminus \{0\}) \quad (2.2)$$

is not essentially self-adjoint and has deficiency indices $(2, 2)$. Consequently, it admits infinitely many self-adjoint extensions. In this paper we will work with the Friedrichs extension of (2.2) which we denote by H_α . In order to define the latter we introduce a function space $W_\alpha^{1,2}(\mathbb{R}^2)$ given by the closure of $C_0^\infty(\mathbb{R}^2 \setminus \{0\})$ with respect to the norm

$$\|u\|_{L^2(\mathbb{R}^2)} + \|(i\nabla + A_{ab})u\|_{L^2(\mathbb{R}^2)}.$$

The quadratic form

$$Q_\alpha[u] = \|(i\nabla + A_{ab})u\|_{L^2(\mathbb{R}^2)}^2 \quad (2.3)$$

with the form domain $W_\alpha^{1,2}(\mathbb{R}^2)$ is then closed and generates a unique non-negative self-adjoint operator H_α in $L^2(\mathbb{R}^2)$. Next we introduce an additional electric potential $V : \mathbb{R}^2 \rightarrow \mathbb{R}$ and consider the Schrödinger operator

$$H = H_\alpha + V \quad \text{in} \quad L^2(\mathbb{R}^2). \quad (2.4)$$

The operators $H_\alpha + V$ and $H_{m+\alpha} + V$ are unitarily equivalent for any $m \in \mathbb{Z}$. We may therefore assume without loss of generality that

$$0 < |\alpha| \leq \frac{1}{2}, \quad \text{which implies} \quad \min_{k \in \mathbb{Z}} |k - \alpha| = |\alpha|.$$

2.1. Notation. Given $s \in \mathbb{R}$ we denote

$$L_s^2(\mathbb{R}^2) = \{u : \|w^s u\|_{L^2(\mathbb{R}^2)} < \infty\}, \quad \|u\|_{0,s} := \|w^s u\|_{L^2(\mathbb{R}^2)},$$

where $w(x) = (1 + |x|^2)^{1/2}$. For $x = (x_1, x_2) \in \mathbb{R}^2$ and $y = (y_1, y_2) \in \mathbb{R}^2$ we will often use the polar coordinates representation

$$x_1 + ix_2 = re^{i\theta}, \quad y_1 + iy_2 = r'e^{i\theta'}, \quad r, r' \geq 0, \quad \theta, \theta' \in [0, 2\pi). \quad (2.5)$$

Given $R > 0$ and a point $x \in \mathbb{R}^2$ we denote by $B(x, R) \subset \mathbb{R}^2$ the open ball with radius R centered in x . By $\|K\|_{\mathcal{B}(s,s')}$ respectively $\|K\|_{\mathcal{B}(\rho,\rho^{-1})}$ we denote the norm of a bounded linear operator K in $\mathcal{B}(s, s')$ respectively $\mathcal{B}(\rho, \rho^{-1})$. The scalar product in a Hilbert space \mathcal{H} will be denoted by $\langle \cdot, \cdot \rangle_{\mathcal{H}}$. Finally, we denote $\mathbb{R}_+ = (0, \infty)$ and $\mathbb{C}_+ = \{z \in \mathbb{C} : \text{Im } z > 0\}$.

3. Main results

3.1. Time decay for e^{-itH_α} . We say that $e^{-itH_\alpha}(x, y)$, $x, y \in \mathbb{R}^2$ is an integral kernel of the operator e^{-itH_α} if

$$(e^{-itH_\alpha} u)(x) = \int_{\mathbb{R}^2} e^{-itH_\alpha}(x, y) u(y) dy \quad (3.1)$$

holds for any $u \in L^2(\mathbb{R}^2)$ with compact support.

Theorem 3.1. *Let $|\alpha| \leq 1/2$. There exists a kernel $e^{-itH_\alpha}(x, y)$ of e^{-itH_α} such that for all $x, y \in \mathbb{R}^2$ it holds*

$$\lim_{t \rightarrow \infty} (it)^{1+|\alpha|} e^{-itH_\alpha}(x, y) = \frac{1}{4\pi\Gamma(1+|\alpha|)} \left(\frac{rr'}{4}\right)^{|\alpha|} \quad \text{if } |\alpha| < 1/2, \quad (3.2)$$

$$\lim_{t \rightarrow \infty} (it)^{\frac{3}{2}} e^{-itH_\alpha}(x, y) = \frac{1}{4\pi\Gamma(3/2)} \left(\frac{rr'}{4}\right)^{\frac{1}{2}} (1 + e^{\mp i(\theta - \theta')}) \quad \text{if } \alpha = \pm 1/2. \quad (3.3)$$

Moreover, there is a constant C such that for all $t > 0$ and all $x, y \in \mathbb{R}^2$

$$|e^{-itH_\alpha}(x, y)| \leq C \min \{t^{-1}, (rr')^{|\alpha|} t^{-1-|\alpha|}\}. \quad (3.4)$$

Remark 3.2. In [FFFP, Thm.1.3 & Cor.1.7] it was shown that

$$\sup_{x, y \in \mathbb{R}^2} |e^{-itH_\alpha}(x, y)| \leq \text{const } t^{-1}.$$

This obviously yields the first part of (3.4). Moreover, the decay rate t^{-1} in the above estimate is sharp. However, the second term on the right hand side of (3.4) shows that for fixed x and y the kernel $e^{-itH_\alpha}(x, y)$ decays faster than t^{-1} .

Inequality (3.4) implies

Corollary 3.3. *There exists $C > 0$ such that for all $t > 0$ we have*

$$\begin{aligned} \|e^{-itH_\alpha} u\|_{0,-s} &\leq C t^{-1-|\alpha|} \|u\|_{0,s} & \forall u \in L_s^2(\mathbb{R}^2), \quad s > 1 + |\alpha|, \\ \|w^{-s} e^{-itH_\alpha} u\|_\infty &\leq C t^{-1-|\alpha|} \|w^s u\|_{L^1(\mathbb{R}^2)} & \forall u \in L_s^1(\mathbb{R}^2), \quad s \geq |\alpha|, \end{aligned}$$

where $L_s^1(\mathbb{R}^2) = \{u : \|w^s u\|_{L^1(\mathbb{R}^2)} < \infty\}$.

Remark 3.4. Improved time decay of semi-groups generated by two-dimensional magnetic Schrödinger operators was recently studied in [Ko, Kr].

3.2. Time decay for $e^{-it(H_\alpha + V)}$. In this case we will introduce another operator norm associated to the weight function $\rho : \mathbb{R}^2 \rightarrow \mathbb{R}$ given by $\rho(x) = \rho(|x|) = e^{|x|^4}$. Let

$$L^2(\mathbb{R}^2, \rho) = \{u : \|\rho^{1/2} u\|_{L^2(\mathbb{R}^2)} < \infty\}, \quad \|u\|_\rho := \|\rho^{1/2} u\|_{L^2(\mathbb{R}^2)}.$$

We denote by $\mathcal{B}(\rho, \rho^{-1})$ the space of bounded linear operators from $L^2(\mathbb{R}^2, \rho)$ to $L^2(\mathbb{R}^2, \rho^{-1})$. Denote by $G_0 \in \mathcal{B}(\rho, \rho^{-1})$ the integral operator with the kernel

$$G_0(x, y) = \frac{1}{4\pi} \sum_{m \in \mathbb{Z}} \frac{e^{im(\theta - \theta')}}{|\alpha + m|} \left(\min \left\{ \frac{r}{r'}, \frac{r'}{r} \right\} \right)^{|\alpha + m|}. \quad (3.5)$$

Note that the series on the right hand side of (3.5) converges for all $x \neq y$. Moreover, from equation (5.4) below it easily follows that $G_0 \in \mathcal{B}(\rho, \rho^{-1})$. As for the potential V , we suppose that it satisfies the following

Assumption 3.5. The function $V : \mathbb{R}^2 \rightarrow \mathbb{R}$ is bounded and compactly supported. Moreover, for any $u \in L^2(\mathbb{R}^2, \rho^{-1})$ it holds

$$u + G_0 V u = 0 \quad \Rightarrow \quad u = 0. \quad (3.6)$$

The motivation for the assumption that V has compact support is twofold. On one hand, it guarantees that $V \in \mathcal{B}(\rho^{-1}, \rho)$ which is needed in (3.6). On the other hand, the compactness of the support of V will play an important role in the proof of absence of positive eigenvalues of the operator $H_\alpha + V$, see Proposition 5.12.

To proceed we note that since V is bounded, the operator $H = H_\alpha + V$ is self-adjoint on the domain of H_α , by Rellich theorem. Let $P_c = P_c(H)$ denote the projection onto the continuous spectral subspace of H . We have

Theorem 3.6. *Let $0 < |\alpha| \leq 1/2$ and let V satisfy assumption 3.5. Denote by $\mathcal{G}_j \in \mathcal{B}(\rho, \rho^{-1})$, $j = 1, 2$, the operators with integral kernels*

$$\begin{aligned} \mathcal{G}_1(x, y) &= \frac{1}{4\pi\Gamma(1+|\alpha|)} \left(\frac{rr'}{4}\right)^{|\alpha|} \\ \mathcal{G}_2(x, y) &= e^{i(\theta' - \theta) \operatorname{sign} \alpha} \frac{\Gamma(1+|\alpha|)}{4\pi\Gamma^2(1+|\alpha - \operatorname{sign} \alpha|)} \left(\frac{rr'}{4}\right)^{|\alpha - \operatorname{sign} \alpha|}. \end{aligned}$$

Then, as $t \rightarrow \infty$

$$e^{-itH} P_c = (it)^{-1-|\alpha|} (1 + G_0 V)^{-1} \mathcal{G}_1 (1 + V G_0)^{-1} + o(t^{-1-|\alpha|}) \quad \text{if } |\alpha| < 1/2, \quad (3.7)$$

$$e^{-itH} P_c = (it)^{-1-|\alpha|} (1 + G_0 V)^{-1} (\mathcal{G}_1 + \mathcal{G}_2) (1 + V G_0)^{-1} + o(t^{-1-|\alpha|}) \quad \text{if } |\alpha| = 1/2 \quad (3.8)$$

in $\mathcal{B}(\rho, \rho^{-1})$. In particular, there exists a constant C such that for all $t > 0$ and all $u \in L^2(\mathbb{R}^2, \rho)$ we have

$$\|e^{-itH} P_c u\|_{\rho^{-1}} \leq C t^{-1-\alpha} \|u\|_\rho. \quad (3.9)$$

Remark 3.7. The reason why we work with the weighted spaces $L^2(\mathbb{R}^2, \rho)$ and not with $L_s^2(\mathbb{R}^2)$ is technical. Our goal is to keep track of the diamagnetic effect on e^{-itH} , i.e. of the improved time decay caused by the magnetic field, observed in the case $V = 0$. Therefore we employ the perturbation approach with H_α as the free operator. This requires a precise knowledge of the behavior of the resolvent of H_α near the threshold of the spectrum. Since we don't have a simple formula for its integral kernel, contrary to the situation without a magnetic field, we work with the power-series (5.2) below. To make sure that this series converges absolutely in $\mathcal{B}(\rho, \rho^{-1})$ we have to require a very fast grow of the weight function ρ . Note that super-polynomially growing weight functions were used to study decay estimates of non-magnetic Schrödinger operators already in [Ra].

Nonetheless, it is reasonable to expect that the claim of Theorem 3.6 remains true also if $L^2(\mathbb{R}^2, \rho)$ and $L^2(\mathbb{R}^2, \rho^{-1})$ are replaced throughout by $L_s^2(\mathbb{R}^2)$ and $L_{-s}^2(\mathbb{R}^2)$ with s large enough.

Remark 3.8. Assumption 3.5 ensures that the operators $1 + G_0 V$ and $1 + V G_0$ are invertible in $\mathcal{B}(\rho^{-1}, \rho^{-1})$ and $\mathcal{B}(\rho, \rho)$ respectively, see the proof of Lemma 5.3 and Remark 5.4 for details. Hence $(1 + G_0 V)^{-1} \mathcal{G}_j (1 + V G_0)^{-1}$ are well defined and belong to $\mathcal{B}(\rho, \rho^{-1})$. Note also that in the case $V = 0$ equations (3.7) and (3.8) agree with the asymptotic (3.2) and (3.3).

Remark 3.9. Our method does not enable us to extend the above results to a more general class of magnetic fields. On the other hand, decay estimates on the magnetic heat semi-group obtained recently in [Ko, Kr] suggest that for a sufficiently smooth magnetic field $B = \operatorname{rot} A$ with a finite total flux α one should be able to observe, in a suitable operator topology, that as $t \rightarrow \infty$

$$\begin{aligned} e^{-it(i\nabla + A)^2} &= \mathcal{O}(t^{-1 - \min_{k \in \mathbb{Z}} |k - \alpha|}) & \text{if } \alpha \notin \mathbb{Z}, \\ e^{-it(i\nabla + A)^2} &= \mathcal{O}(t^{-1} \log^{-2} t) & \text{if } \alpha \in \mathbb{Z}. \end{aligned}$$

This question remains open.

4. Partial wave decomposition

The quadratic form associated to H_α in polar coordinates (r, θ) reads as follows:

$$Q_\alpha[u] = \int_0^\infty \int_0^{2\pi} (|\partial_r u|^2 + r^{-2} |i \partial_\theta u + \alpha u|^2) r dr d\theta, \quad u \in W_\alpha^{1,2}(\mathbb{R}^2). \quad (4.1)$$

By expanding a given function $u \in L^2(\mathbb{R}_+ \times (0, 2\pi))$ into a Fourier series with respect to the basis $\{e^{im\theta}\}_{m \in \mathbb{Z}}$ of $L^2((0, 2\pi))$, we obtain a direct sum decomposition

$$H_\alpha = \sum_{m \in \mathbb{Z}} \oplus (h_m \otimes \text{id}) \Pi_m, \quad (4.2)$$

where h_m are operators generated by the closures, in $L^2(\mathbb{R}_+, r dr)$, of the quadratic forms

$$Q_m[f] = \int_0^\infty \left(|f'|^2 + \frac{(\alpha + m)^2}{r^2} |f|^2 \right) r dr \quad (4.3)$$

defined initially on $C_0^\infty(\mathbb{R}_+)$, and Π_m is the projector acting as

$$(\Pi_m u)(r, \theta) = \frac{1}{2\pi} \int_0^{2\pi} e^{im(\theta - \theta')} u(r, \theta') d\theta'.$$

5. Resolvent estimates

In order to prove Theorem 3.6 we will follow the strategy developed in [JK]. This requires precise estimates on the resolvent of $(H - z)^{-1}$ and its derivatives with respect to z . Such estimates are the main objects of our interest in this section.

5.1. Low energy behavior. Let us denote by $R_0(\alpha, z) = (H_\alpha - z)^{-1}$ the resolvent of the free operator H_α . In order to study the behavior of $R_0(\alpha, \lambda)$ for $\lambda \rightarrow 0$, $\lambda \in (0, \infty)$, we first consider the resolvent kernels of the one-dimensional operators h_m associated with quadratic form (4.3). From [Ko, Sect.5] it follows that for $\lambda \in (0, \infty)$ and $r < r'$

$$(h_m - \lambda)^{-1}(r, r') = \frac{\pi i}{2} J_{|m+\alpha|}(r\sqrt{\lambda}) (J_{|m+\alpha|}(r'\sqrt{\lambda}) + i Y_{|m+\alpha|}(r'\sqrt{\lambda})),$$

where J_μ and Y_μ are the Bessel functions of the first and second kind respectively. When $r' \leq r$, then we switch r' and r in the above formula. In the sequel we will assume for definiteness that $r < r'$. By (4.2) and (2.5) we get

$$R_0(\alpha, \lambda, x, y) = \frac{i}{4} \sum_{m \in \mathbb{Z}} J_{|\alpha+m|}(r\sqrt{\lambda}) (J_{|\alpha+m|}(r'\sqrt{\lambda}) + i Y_{|\alpha+m|}(r'\sqrt{\lambda})) e^{im(\theta - \theta')} \quad (5.1)$$

Using [AS, Eqs. 9.1.2, 9.1.10] and the well-known properties of the Gamma function, namely

$$\Gamma(z+1) = z \Gamma(z), \quad \Gamma(z) \Gamma(1-z) = \frac{\pi}{\sin(\pi z)},$$

we can further rewrite (5.1) in the power-series in λ as follows:

$$\begin{aligned}
R_0(\alpha, \lambda, x, y) &= G_0(x, y) + G_1(x, y) \lambda^{|\alpha|} + G_2(x, y) \lambda^{|\alpha - \text{sign } \alpha|} \\
&+ \frac{1}{4\pi} \sum_{m \in \mathbb{Z}} e^{im(\theta - \theta')} \left(\frac{r}{r'}\right)^{|\alpha+m|} \sum_{k, n \geq 0, k+n > 0} \frac{\left(\frac{1}{4} r'^2\right)^k \left(-\frac{1}{4} r^2\right)^n \lambda^{k+n}}{n! k! (n + |\alpha + m|) \cdots |\alpha + m| \cdots (|\alpha + m| - k)} \\
&+ \frac{i - \cot(\pi|\alpha|)}{4^{1+|\alpha|}} \sum_{k, n \geq 0, k+n > 0} \frac{\left(-\frac{1}{4} r'^2\right)^k \left(-\frac{1}{4} r^2\right)^n (rr')^{|\alpha|} \lambda^{k+n+|\alpha|}}{\Gamma(n + |\alpha| + 1) \Gamma(k + |\alpha| + 1)} \\
&+ \sum_{m \neq 0} e^{im(\theta - \theta')} \frac{i - \cot(\pi|\alpha + m|)}{4^{1+|\alpha+m|}} \sum_{k, n \geq 0, k+n > 0} \frac{\left(-\frac{1}{4} r'^2\right)^k \left(-\frac{1}{4} r^2\right)^n (rr')^{|\alpha+m|} \lambda^{k+n+|\alpha+m|}}{\Gamma(n + |\alpha + m| + 1) \Gamma(k + |\alpha + m| + 1)},
\end{aligned} \tag{5.2}$$

where $G_0(x, y)$ is given by (3.5) and

$$\begin{aligned}
G_1(x, y) &= \frac{i - \cot(\pi|\alpha|)}{4\Gamma^2(1 + |\alpha|)} \left(\frac{rr'}{4}\right)^{|\alpha|}, \\
G_2(x, y) &= e^{i(\theta' - \theta) \text{sign } \alpha} \frac{i - \cot(\pi|\alpha - \text{sign } \alpha|)}{4\Gamma^2(1 + |\alpha - \text{sign } \alpha|)} \left(\frac{rr'}{4}\right)^{|\alpha - \text{sign } \alpha|}.
\end{aligned}$$

Denote by G_j , $j = 1, 2$, the respective integral operators generated by the kernels $G_j(x, y)$.

Lemma 5.1. *The series of integral operators (5.2) converges absolutely in $\mathcal{B}(\rho, \rho^{-1})$ and uniformly in λ on compacts of $[0, \infty)$. In particular, we have*

$$R_0(\alpha, \lambda) = G_0 + G_1 \lambda^{|\alpha|} + G_2 \lambda^{|\alpha - \text{sign } \alpha|} + o(\lambda^{|\alpha - \text{sign } \alpha|}) \quad \text{in } \mathcal{B}(\rho, \rho^{-1}). \tag{5.3}$$

as $\lambda \rightarrow 0$. Moreover, the operator $R_0(\alpha, \lambda)$ generated by the integral kernel (5.2) can be differentiated in λ on $(0, 1)$ any number of times in the norm of $\mathcal{B}(\rho, \rho^{-1})$.

Proof. If K is an integral operator from $L^2(\mathbb{R}^2, \rho)$ to $L^2(\mathbb{R}^2, \rho^{-1})$ with an integral kernel $k(x, y)$, then

$$\|K\|_{\mathcal{B}(\rho, \rho^{-1})} \leq \|K\|_{HS} = \left(\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} |k(x, y)|^2 \rho^{-1}(x) \rho^{-1}(y) dx dy \right)^{1/2}, \tag{5.4}$$

where $\|\cdot\|_{HS}$ denotes the Hilbert-Schmidt norm. It is thus easily verified that the series of operators on the right hand side of (5.2) converges absolutely in $\mathcal{B}(\rho, \rho^{-1})$ for any $\lambda \geq 0$ and that the convergence is uniform in λ on any compact interval of $[0, \infty)$. The same argument applies to $\frac{d^r}{d\lambda^r} R_0(\alpha, \lambda)$ for any $r \in \mathbb{N}$. \square

Corollary 5.2. *Let $R'_0(\alpha, \lambda)$ and $R''_0(\alpha, \lambda)$ denote first and second derivative of $R_0(\alpha, \lambda)$ with respect to λ in the norm of $\mathcal{B}(\rho, \rho^{-1})$. Then, as $\lambda \rightarrow 0$, we have*

$$R'_0(\alpha, \lambda) = |\alpha| G_1 \lambda^{|\alpha|-1} + o(\lambda^{|\alpha|-1}), \quad R''_0(\alpha, \lambda) = |\alpha|(|\alpha| - 1) G_1 \lambda^{|\alpha|-2} + o(\lambda^{|\alpha|-2}). \tag{5.5}$$

in $\mathcal{B}(\rho, \rho^{-1})$.

Proof. This follows from Lemma 5.1. \square

The perturbed resolvent. Next we consider the full resolvent $R(\alpha, z) = (H - z)^{-1}$ of the operator $H = H_\alpha + V$ in $L^2(\mathbb{R}^2)$.

Lemma 5.3. *Under Assumption 3.5 we have*

$$(1 + R_0(\alpha, \lambda) V)^{-1} = (1 + G_0 V)^{-1} - (1 + G_0 V)^{-1} G_1 V (1 + G_0 V)^{-1} \lambda^{|\alpha|} + o(\lambda^{|\alpha|}) \tag{5.6}$$

as $\lambda \rightarrow 0$ in $\mathcal{B}(\rho^{-1}, \rho^{-1})$.

Proof. Both operators G_0V and G_1V are Hilbert-Schmidt, and therefore compact, from $L^2(\mathbb{R}^2, \rho^{-1})$ to $L^2(\mathbb{R}^2, \rho^{-1})$. Hence by (3.6) the operator $1 + G_0V$ is invertible in $\mathcal{B}(\rho^{-1}, \rho^{-1})$. On the other hand, in view of (5.3) we have

$$1 + R_0(\alpha, \lambda)V = 1 + G_0V + G_1V\lambda^{|\alpha|} + o(\lambda^{|\alpha|}) \quad (5.7)$$

in $\mathcal{B}(\rho^{-1}, \rho^{-1})$. Here we have used the fact that $V \in \mathcal{B}(\rho^{-1}, \rho)$. It follows that for λ small enough the operator $1 + R_0(\alpha, \lambda)V$ can be inverted, and with the help of the Neumann series and we arrive at (5.6). \square

Remark 5.4. The operator VG_0 is compact from $L^2(\mathbb{R}^2, \rho)$ to $L^2(\mathbb{R}^2, \rho)$. Hence by equation (3.6) and duality $1 + VG_0$ is invertible in $\mathcal{B}(\rho, \rho)$.

Lemma 5.5. *Under Assumption 3.5 we have*

$$R(\alpha, \lambda) = T_0 + T_1\lambda^{|\alpha|} + o(\lambda^{|\alpha|}) \quad (5.8)$$

as $\lambda \rightarrow 0$ in $\mathcal{B}(\rho, \rho^{-1})$, where

$$T_0 = (1 + G_0V)^{-1}G_0, \quad T_1 = (1 + G_0V)^{-1}G_1(1 + VG_0)^{-1}. \quad (5.9)$$

Proof. Since $1 + R_0(\alpha, \lambda)V$ is invertible in $\mathcal{B}(\rho^{-1}, \rho^{-1})$ for λ small enough, the resolvent equation yields

$$R(\alpha, \lambda) = (1 + R_0(\alpha, \lambda)V)^{-1}R_0(\alpha, \lambda), \quad (5.10)$$

which in combination with (5.3) and (5.6) gives equation (5.8) with

$$T_1 = (1 + G_0V)^{-1}G_1 - (1 + G_0V)^{-1}G_1V(1 + G_0V)^{-1}G_0$$

in $\mathcal{B}(\rho, \rho^{-1})$. To simplify the above expression for T_1 let us consider $u \in L^2(\mathbb{R}^2, \rho)$ and denote $f = G_0u \in L^2(\mathbb{R}^2, \rho^{-1})$. Since $(1 + G_0V)^{-1}f = g$ if and only if $f = g + G_0Vg$, and $(1 + VG_0)^{-1}Vf = Vg$ if and only if $Vf = Vg + VG_0Vg$, we find out that $V(1 + G_0V)^{-1}G_0u = (1 + VG_0)^{-1}VG_0u$. Hence the identity

$$V(1 + G_0V)^{-1}G_0 = (1 + VG_0)^{-1}VG_0 \quad (5.11)$$

holds on $L^2(\mathbb{R}^2, \rho)$, which implies that

$$T_1 = (1 + G_0V)^{-1}G_1(1 + VG_0)^{-1}.$$

\square

Corollary 5.6. *As $\lambda \rightarrow 0$,*

$$R''(\alpha, \lambda) = |\alpha|(|\alpha| - 1)(1 - T_0V)G_1(1 - T_0V)\lambda^{|\alpha|-2} + o(\lambda^{|\alpha|-2}) \quad \text{in } \mathcal{B}(\rho, \rho^{-1}). \quad (5.12)$$

Proof. We use the identities

$$R'(\alpha, \lambda) = (1 - R(\alpha, \lambda)V)R'_0(\alpha, \lambda)(1 - VR(\alpha, \lambda)) \quad (5.13)$$

$$R''(\alpha, \lambda) = (1 - R(\alpha, \lambda)V)R''_0(\alpha, \lambda)(1 - VR(\alpha, \lambda)) - 2R'(\alpha, \lambda)VR'_0(\alpha, \lambda)(1 - VR(\alpha, \lambda)), \quad (5.14)$$

which hold in $\mathcal{B}(\rho, \rho^{-1})$ in view of the resolvent equation $R = R_0 - R_0VR$. The claim now follows from Corollary 5.2 and Lemma 5.5. \square

5.2. High energy behavior. Here we will use the operator norms in the weighted spaces $L_s^2(\mathbb{R}^2)$. When $s = 0$ we write $\|u\|_0$ instead of $\|u\|_{0,0}$. We denote by $\mathcal{B}(s, s')$ the set of bounded operators from $L_{s'}^2(\mathbb{R}^2)$ to $L_s^2(\mathbb{R}^2)$. Obviously, for any $K \in \mathcal{B}(s, s')$ we have

$$p' \leq s' \leq s \leq p \quad \Rightarrow \quad \|K\|_{\mathcal{B}(p, p')} \leq \|K\|_{\mathcal{B}(s, s')}. \quad (5.15)$$

According to [IT, Prop. 7.3], the operator $R_0(\alpha, \lambda)$ belongs to $\mathcal{B}(s, -s)$ for any $s > 1/2$, and the limit

$$\lim_{\varepsilon \rightarrow 0+} R_0(\alpha, \lambda + i\varepsilon) = R_0(\alpha, \lambda), \quad \text{in } \mathcal{B}(s, -s) \quad (5.16)$$

is attained locally uniformly in λ on $(0, \infty)$.

Lemma 5.7. *If $s > 1/2$, then $R_0(\alpha, \lambda)$ is continuous in λ on $(0, \infty)$ in the norm of $\mathcal{B}(s, -s)$.*

Proof. First we show that for any s with $s \leq 1$ and any $z \in \mathbb{C}$ with $\text{Im } z \neq 0$ we have $R_0(\alpha, z) \in \mathcal{B}(s, s)$. Given $u \in L_s^2(\mathbb{R}^2)$, we have

$$\begin{aligned} \|R_0(\alpha, z) u\|_{0,s} &\leq \|R_0(\alpha, z) w^s u\|_0 + \|[R_0(\alpha, z), w^s] u\|_0 \\ &= \|R_0(\alpha, z) w^s u\|_0 + \|[R_0(\alpha, z) [H_\alpha, w^s] R_0(\alpha, z) u]\|_0. \end{aligned} \quad (5.17)$$

In the polar coordinates H_α acts as

$$H_\alpha = -\partial_r^2 - \frac{1}{r} \partial_r + \frac{1}{r^2} (i\partial_\theta + \alpha)^2.$$

Hence when calculated on functions from $C_0^\infty(\mathbb{R}^2 \setminus \{0\})$ the commutator $[H_\alpha, w^s]$ reads as

$$[H_\alpha, w^s] = -s w(r)^{s-2} (sr^2 + 2) - 2s r w(r)^{s-2} \partial_r,$$

where $r = |x|$. Since $s \leq 1$, the first term on the right hand side is bounded. Moreover, from (4.1) it follows that for every $f \in L^2(\mathbb{R}^2)$

$$\|\partial_r R_0(\alpha, z) f\|_{L^2(\mathbb{R}^2)}^2 \leq |\langle f, R_0(\alpha, z) f \rangle_{L^2(\mathbb{R}^2)}| + |z| \|R_0(\alpha, z) f\|_{L^2(\mathbb{R}^2)}^2.$$

This in combination with (5.17) and the fact that

$$\|R_0(\alpha, z)\|_{\mathcal{B}(0,0)} \leq \frac{1}{|\text{Im } z|} \quad (5.18)$$

gives

$$\|R_0(\alpha, z) u\|_{0,s} \leq C \left(\frac{1}{|\text{Im } z|} + \frac{|z|}{|\text{Im } z|^2} \right) \|u\|_{0,s}. \quad (5.19)$$

Now, let $\lambda, \lambda' \in \mathbb{R}$ and let $\varepsilon > 0$. Then by the resolvent equation

$$\|(R_0(\alpha, \lambda + i\varepsilon) - R_0(\alpha, \lambda' + i\varepsilon)) u\|_{0,s} = |\lambda - \lambda'| \|R_0(\alpha, \lambda + i\varepsilon) R_0(\alpha, \lambda' + i\varepsilon) u\|_{0,s}. \quad (5.20)$$

Hence in view of (5.19), for ε small enough

$$\begin{aligned} \|(R_0(\alpha, \lambda) - R_0(\alpha, \lambda')) u\|_{0,-s} &\leq \|R_0(\alpha, \lambda) - R_0(\alpha, \lambda + i\varepsilon)\|_{\mathcal{B}(s, -s)} \|u\|_{0,s} \\ &\quad + \|R_0(\alpha, \lambda') - R_0(\alpha, \lambda' + i\varepsilon)\|_{\mathcal{B}(s, -s)} \|u\|_{0,s} + \mathcal{O}(\varepsilon^{-4}) |\lambda - \lambda'| (1 + |\lambda|)(1 + |\lambda'|) \|u\|_{0,s} \end{aligned}$$

Since the first two terms on the right hand side converge to zero as $\varepsilon \rightarrow 0$ locally uniformly in λ respectively λ' , see (5.16), this proves the continuity of $R_0(\alpha, \lambda)$ in $\lambda \in (0, \infty)$ for $s \in (1/2, 1]$. By (5.15) the claim holds for all $s > 1/2$. \square

Lemma 5.8. *For any $\varepsilon > 0$ there exists C_ε such that for all $s, s' \geq 1/2 + \varepsilon$ it holds*

$$\|R_0(\alpha, \lambda)\|_{\mathcal{B}(s, -s')} \leq C_\varepsilon \lambda^{-\frac{1}{2} + \varepsilon} \quad \forall \lambda \geq 1. \quad (5.21)$$

Proof. Let $s = 1/2 + \varepsilon$ and let $u \in L_s^2(\mathbb{R}^2)$. We define $u_\lambda(x) = u(x/\sqrt{\lambda})$. Since $R_0(\alpha, \lambda, x, y) = R_0(\alpha, 1, \sqrt{\lambda}x, \sqrt{\lambda}y)$, see equation (5.1), a simple change of variables gives

$$\|R_0(\alpha, \lambda) u\|_{0,-s}^2 \leq \lambda^{-3+s} \|R_0(\alpha, 1)\|_{\mathcal{B}(s,-s)}^2 \|u_\lambda\|_{0,s}^2.$$

On the other hand, $\|u_\lambda\|_{0,s}^2 \leq \lambda^{1+s} \|u\|_{0,s}^2$. Hence

$$\|R_0(\alpha, \lambda) u\|_{0,-s}^2 \leq \lambda^{-1+2\varepsilon} \|R_0(\alpha, 1)\|_{\mathcal{B}(s,-s)}^2 \|u\|_{0,s}^2.$$

In view of (5.15) this completes the proof. \square

In order to obtain suitable estimates on the derivatives of $R_0(\alpha, \lambda)$, we need the following technical result.

Lemma 5.9. *Let $T = (i\nabla + A_{ab}) \cdot x$. Assume that $s, s' > 5/2$. Then for any $\varepsilon > 0$ we have*

$$[iT, R_0(\alpha, \lambda)] = \mathcal{O}(\lambda^\varepsilon) \quad \text{in } \mathcal{B}(s, -s') \quad \text{as } \lambda \rightarrow \infty.$$

Proof. We introduce the notation $D_j = -\partial_j + iA_j$, $j = 1, 2$ so that $iT = D_1x_1 + D_2x_2$. We observe that $[D_j, x_j] = -1$. Hence in view of Lemma 5.8 in order to show that $\|T R_0(\alpha, \lambda)\|_{\mathcal{B}(s,-s')} = \mathcal{O}(\lambda^\varepsilon)$, it suffices to prove that for any $p > 1/2$ and any $u \in L_p^2(\mathbb{R}^2)$

$$\begin{aligned} \|w^{-\frac{s'}{2}}(x_1 D_1 + x_2 D_2) R_0(\alpha, \lambda) u\|_0^2 &\leq 2 \|w^{1-\frac{s'}{2}} D_1 R_0(\alpha, \lambda) u\|_0^2 + 2 \|w^{1-\frac{s'}{2}} D_2 R_0(\alpha, \lambda) u\|_0^2 \\ &=: 2 X_1^2(\lambda) + 2 X_2^2(\lambda) = \mathcal{O}(\lambda^{2\varepsilon}) \|u\|_{0,p}^2 \end{aligned} \quad (5.22)$$

as $\lambda \rightarrow \infty$. Note that $w^{\frac{s'}{2}-1} [D_j, w^{1-\frac{s'}{2}}] = -w^{\frac{s'}{2}-1} \partial_j w^{1-\frac{s'}{2}}$ is a bounded function for $j = 1, 2$. Hence by the Cauchy-Schwarz inequality and (5.21) we get

$$\begin{aligned} X_j^2(\lambda) &= \langle w^{1-\frac{s'}{2}} R_0(\alpha, \lambda) u, [D_j^*, w^{1-\frac{s'}{2}}] D_j R_0(\alpha, \lambda) u \rangle_{L^2(\mathbb{R}^2)} + \\ &+ \langle [w^{1-\frac{s'}{2}}, D_j] R_0(\alpha, \lambda) u, w^{1-\frac{s'}{2}} D_j R_0(\alpha, \lambda) u \rangle_{L^2(\mathbb{R}^2)} + \langle w^{1-\frac{s'}{2}} R_0(\alpha, \lambda) u, w^{1-\frac{s'}{2}} D_j^* D_j R_0(\alpha, \lambda) u \rangle_{L^2(\mathbb{R}^2)} \\ &\leq 2 C_\varepsilon \lambda^{-\frac{1}{2}+\varepsilon} X_j(\lambda) \|u\|_{0,p} + \langle w^{1-\frac{s'}{2}} R_0(\alpha, \lambda) u, w^{1-\frac{s'}{2}} D_j^* D_j R_0(\alpha, \lambda) u \rangle_{L^2(\mathbb{R}^2)}. \end{aligned}$$

In the application of (5.21) we used the assumption $s' - 2 > 1/2$. Now from the identity

$$\sum_{j=1}^2 D_j^* D_j R_0(\alpha, \lambda) = H_\alpha R_0(\alpha, \lambda) = 1 + \lambda R_0(\alpha, \lambda)$$

and equation (5.21) we obtain

$$X_1^2(\lambda) + X_2^2(\lambda) \leq 4 C_\varepsilon \lambda^{-\frac{1}{2}+\varepsilon} (X_1(\lambda) + X_2(\lambda)) \|u\|_{0,p} + C_\varepsilon \lambda^{-\frac{1}{2}+\varepsilon} \|u\|_{0,p}^2 + C_\varepsilon^2 \lambda^{2\varepsilon} \|u\|_{0,p}^2.$$

This implies (5.22). To prove that $\|R_0(\alpha, \lambda) T\|_{\mathcal{B}(s,-s')} = \mathcal{O}(\lambda^\varepsilon)$, let $u \in C_0^\infty(\mathbb{R}^2 \setminus \{0\})$ and write $v_j = x_j u$. From (5.22) applied to v_j with $p = s - 2$ we deduce that

$$\|w^{-\frac{s'}{2}}(D_1 R_0(\alpha, \lambda) v_1 + D_2 R_0(\alpha, \lambda) v_2)\|_0^2 \leq 2 \sum_{j=1}^2 \|w^{-\frac{s'}{2}} D_j R_0(\alpha, \lambda) v_j\|_0^2 = \mathcal{O}(\lambda^{2\varepsilon}) \|u\|_{0,s}^2 \quad (5.23)$$

as $\lambda \rightarrow \infty$. Since the operators D_j commute with $R_0(\alpha, \lambda)$ on $C_0^\infty(\mathbb{R}^2 \setminus \{0\})$, inequality (5.23) implies that

$$\|R_0(\alpha, \lambda) T u\|_{-s'}^2 = \mathcal{O}(\lambda^{2\varepsilon}) \|u\|_{0,s}^2 \quad \text{as } \lambda \rightarrow \infty$$

for all $u \in C_0^\infty(\mathbb{R}^2 \setminus \{0\})$. By density we conclude that $\|R_0(\alpha, \lambda) T\|_{\mathcal{B}(s,-s')} = \mathcal{O}(\lambda^\varepsilon)$. \square

Lemma 5.10. *Let $s, s' > 5/2$. Let $R_0^{(k)}(\alpha, \lambda)$ denote the k -th derivative of $R_0(\alpha, \lambda)$ with respect to λ in $\mathcal{B}(s, -s')$. Then for any $\varepsilon > 0$ there exists $C_{\varepsilon,k}$ such that*

$$\|R_0^{(k)}(\alpha, \lambda)\|_{\mathcal{B}(s,-s')} \leq C_{\varepsilon,k} \lambda^{-\frac{k+1}{2}+\varepsilon} \quad \forall \lambda \geq 1, \quad k = 1, 2. \quad (5.24)$$

Proof. Let $T = (i\nabla + A) \cdot x$ be the operator introduced in Lemma 5.9. Then the commutator $[iT, H_\alpha]$ calculated on $C_0^\infty(\mathbb{R}^2 \setminus \{0\})$ gives

$$[iT, H_\alpha] = 2H_\alpha.$$

This together with the first resolvent equation $R_0(\alpha, z) - R_0(\alpha, \zeta) = (z - \zeta)R_0(\alpha, z)R_0(\alpha, \zeta)$, $z, \zeta \in \mathbb{C}_+$, implies the identity

$$[iT, R_0(\alpha, z)] = R_0(\alpha, z)[iT, H_\alpha]R_0(\alpha, z) = 2R_0(\alpha, z) + 2zR'_0(\alpha, z). \quad (5.25)$$

On the other hand, from Lemma 5.9 it follows that $[iT, R_0(\alpha, \lambda)] \in \mathcal{B}(s, -s')$ for $\lambda \geq 1$. Equation (5.25), considered in $\mathcal{B}(s, -s')$, thus can be thus extended to $z = \lambda \in \mathbb{R}$, $\lambda \geq 1$. This shows that $R'_0(\alpha, \lambda)$ is continuous in λ on $(1, \infty)$ with respect to the norm of $\mathcal{B}(s, -s')$ and in view of Lemma 5.9

$$R'_0(\alpha, \lambda) = \mathcal{O}(|\lambda|^{-1+\varepsilon}) \quad \text{in } \mathcal{B}(s, -s'), \quad \lambda \rightarrow \infty. \quad (5.26)$$

Hence equation (5.24) is proven for $k = 1$. To prove it for $k = 2$ we consider the double commutator

$$[x, [x, R_0(\alpha, z)]] = |x|^2 R_0(\alpha, z) - 2 \sum_{j=1}^2 x_j R_0(\alpha, z) x_j + R_0(\alpha, z) |x|^2$$

and use the identity

$$z R''_0(\alpha, z) = -3R'_0(\alpha, z) - \frac{1}{4} [x, [x, R_0(\alpha, z)]], \quad (5.27)$$

which follows by a direct calculation in the same way as equation (5.25). By using the fact that

$$x \in \mathcal{B}(s, s-1) \quad \forall s \in \mathbb{R}, \quad (5.28)$$

we extend equation (5.27), considered in $\mathcal{B}(s, -s')$, to $z = \lambda \in \mathbb{R}$, $\lambda \geq 1$. From (5.24) for $k = 1$, (5.26), and (5.21) we thus arrive at

$$R''_0(\alpha, \lambda) = \mathcal{O}(|\lambda|^{-\frac{3}{2}+\varepsilon}) \quad \text{in } \mathcal{B}(s, -s'), \quad \lambda \rightarrow \infty.$$

The claim now follows. \square

Corollary 5.11. *For any $\varepsilon > 0$ and any $k = 0, 1, 2$ there exist constants $C_{\varepsilon, k}$ such that*

$$\|R_0^{(k)}(\alpha, \lambda)\|_{\mathcal{B}(\rho, \rho^{-1})} \leq C_{\varepsilon, k} \lambda^{-\frac{k+1}{2}+\varepsilon} \quad \forall \lambda \geq 1. \quad (5.29)$$

Proof. Since for any $S > 0$ there exists C_s such that

$$\|R_0(\alpha, \lambda)\|_{\mathcal{B}(\rho, \rho^{-1})} \leq C_s \|R_0(\alpha, \lambda)\|_{\mathcal{B}(s, -s)},$$

the claim follows from Lemmata 5.8, 5.10. \square

Proposition 5.12. *Suppose that Assumption 3.5 is satisfied. Then, for any $k = 0, 1, 2$ we have $R^{(k)}(\alpha, \lambda) \in \mathcal{B}(\rho, \rho^{-1})$ and*

$$\|R^{(k)}(\alpha, \lambda)\|_{\mathcal{B}(\rho, \rho^{-1})} = \mathcal{O}(\lambda^{-\frac{k+1}{2}+0}) \quad \lambda \rightarrow \infty. \quad (5.30)$$

Proof. Let us first show that the operator $H = H_\alpha + V$ has no positive eigenvalues. To this end assume that there exists $\lambda > 0$ such that $Hu = \lambda u$. Since V is compactly supported, there exists $R > 0$ such that $V = 0$ outside the ball $B(0, R)$. We can thus use decomposition (4.2) to obtain

$$h_m f_m(r) = \lambda f_m(r) \quad \forall m \in \mathbb{Z}, \quad \forall r > R, \quad (5.31)$$

where

$$f_m(r) = \frac{1}{2\pi} \langle e^{im\theta}, u(r, \theta) \rangle_{L^2(0, 2\pi)}$$

and h_m is the operator associated with quadratic form Q_m given by (4.3). A direct calculation now shows that f_m is a linear combination of the Bessel functions $J_{|m+\alpha|}(r\sqrt{\lambda})$ and $Y_{|m+\alpha|}(r\sqrt{\lambda})$. Since

any non-trivial linear combination of these functions lies outside $L^2((R, \infty), r dr)$, see [AS, Eqs.9.2.1-2], we conclude that $f_m = 0$ on (R, ∞) . By the unique continuation principle we then have $f_m = 0$ on \mathbb{R}_+ for all m and therefore $u = 0$. Hence λ is not an eigenvalue.

Next, since $R_0(\alpha, \lambda)$ is Hilbert-Schmidt, and therefore compact, from $L^2(\mathbb{R}^2, \rho)$ to $L^2(\mathbb{R}^2, \rho^{-1})$, and $V \in \mathcal{B}(\rho^{-1}, \rho)$, we observe that $VR_0(\alpha, \lambda)$ is compact from $L^2(\mathbb{R}^2, \rho)$ to $L^2(\mathbb{R}^2, \rho)$. Moreover the fact that H has no positive eigenvalues implies that -1 is not an eigenvalue of $VR_0(\alpha, \lambda)$ in $\mathcal{B}(\rho, \rho)$ for any $\lambda \in (0, \infty)$. Hence the operator $1 + VR_0(\alpha, \lambda)$ is invertible in $\mathcal{B}(\rho, \rho)$ and its inverse is continuous with respect to λ in the norm of $\mathcal{B}(\rho, \rho)$. The latter follows from the continuity of $VR_0(\alpha, \lambda)$ in $\mathcal{B}(\rho, \rho)$. Now equation (5.10) implies that $R(\alpha, \lambda) \in \mathcal{B}(\rho, \rho^{-1})$ and that $R(\alpha, \lambda)$ is continuous in $\lambda \in (0, \infty)$.

Finally, since $\|VR_0(\alpha, \lambda)\|_{\mathcal{B}(\rho, \rho)} \rightarrow 0$ as $\lambda \rightarrow \infty$, by Corollary 5.11, we conclude that $(1 + VR_0(\alpha, \lambda))^{-1}$ is uniformly bounded for $\lambda \rightarrow \infty$ in the norm of $\mathcal{B}(\rho, \rho)$. In view of (5.10) and Corollary 5.11 we thus get

$$\|R(\alpha, \lambda)\|_{\mathcal{B}(\rho, \rho^{-1})} = \mathcal{O}(\lambda^{-\frac{1}{2}+0}) \quad \lambda \rightarrow \infty.$$

This proves (5.30) for $k = 0$. The proof for $k = 1$ and $k = 2$ now follows from (5.30) with $k = 0$, identities (5.13), (5.14) and Corollary 5.11. \square

6. Time decay

6.1. Proof of Theorem 3.6. For the sake of brevity we will prove the statements of Theorem 3.6 only for $|\alpha| < 1/2$. The proof in the case $|\alpha| = 1/2$ is completely analogous. By the spectral theorem and the Stone formula we have

$$e^{-itH} P_c = \int_0^\infty e^{-it\lambda} E(\alpha, \lambda) d\lambda, \quad (6.1)$$

where

$$E(\alpha, \lambda) = \frac{1}{\pi} \operatorname{Im} R(\alpha, \lambda). \quad (6.2)$$

Let $\chi \in C^\infty(0, \infty)$ be such that $\chi(x) = 0$ for all x large enough and $\chi(x) = 1$ in a neighborhood of 0. Since $E''(\alpha, \lambda) \in L^1((\delta, \infty); \mathcal{B}(\rho, \rho^{-1}))$ for any $\delta > 0$ in view of Proposition 5.12, we can apply [JK, Lemma 10.1], see Lemma A.1 in Appendix A, to obtain

$$\int_0^\infty e^{-it\lambda} (1 - \chi(\lambda)) E(\alpha, \lambda) d\lambda = o(t^{-2}) \quad \text{as } t \rightarrow \infty \quad (6.3)$$

in $\mathcal{B}(\rho, \rho^{-1})$. On the other hand, for $\lambda \rightarrow 0$ we have by (6.2) and Lemma 5.5

$$E(\alpha, \lambda) = E_1 \lambda^{|\alpha|} + E_2(\alpha, \lambda), \quad E_2(\alpha, \lambda) = o(\lambda^{|\alpha|}),$$

where

$$E_1 = \frac{1}{\Gamma(1 + |\alpha|)} (1 + G_0 V)^{-1} \mathcal{G}_1 (1 + V G_0)^{-1}.$$

Moreover, by Corollary 5.6 we know that $E_2''(\alpha, \lambda) = o(\lambda^{|\alpha|-2})$ as $\lambda \rightarrow 0$. Hence from [JK, Lemma 10.2], see Lemma A.1 in Appendix A, applied to $\chi(\lambda) E_2(\alpha, \lambda)$ it follows that

$$\int_0^\infty e^{-it\lambda} \chi(\lambda) E_2(\alpha, \lambda) d\lambda = o(t^{-1-|\alpha|}) \quad t \rightarrow \infty.$$

Finally, since $\chi(\lambda) = 1$ in a vicinity of zero, the Erdelyi's lemma about asymptotic expansion of Fourier integrals, see [Erd2] or [Z, p. 639], gives

$$\int_0^\infty e^{-it\lambda} \lambda^{|\alpha|} \chi(\lambda) d\lambda = (it)^{-1-|\alpha|} \Gamma(1 + |\alpha|) + o(t^{-1-|\alpha|}) \quad t \rightarrow \infty.$$

Summing up, we have

$$e^{-itH} P_c = (it)^{-1-|\alpha|} \Gamma(1 + |\alpha|) E_1 + o(t^{-1-|\alpha|})$$

in $\mathcal{B}(\rho, \rho^{-1})$. This completes the proof of (3.7). Equation (3.8) follows in the same way with E_1 replaced by

$$\frac{1}{\Gamma(1+|\alpha|)} (1+G_0V)^{-1} (\mathcal{G}_1+\mathcal{G}_2) (1+VG_0)^{-1}.$$

Estimate (3.9) is now immediate since $\|e^{-itH} P_c u\|_{\rho^{-1}} \leq \|u\|_\rho$ for all $t > 0$.

Remark 6.1. In principle it would be possible to extend the above analysis and obtain higher order terms in the asymptotic expansion for $e^{-itH} P_c$ as $t \rightarrow \infty$.

6.2. Proof of Theorem 3.1. We first establish an explicit formula for $e^{-itH_\alpha}(x, y)$. This has been done already in [FFFP]. For the sake of completeness we provide an alternative derivation. To start with we consider the one-dimensional operators h_m in $L^2(\mathbb{R}_+, r dr)$ associated with the closure of the quadratic form Q_m given by equation (4.3).

Lemma 6.2. *Let $f \in L^2(\mathbb{R}_+, r dr)$ be compactly supported. Then for all $t > 0$ we have*

$$(e^{-it h_m} f)(r) = \frac{1}{2it} \int_0^\infty I_{|m+\alpha|} \left(\frac{rr'}{2it} \right) e^{-\frac{r^2+r'^2}{4it}} f(r') r' dr'. \quad (6.4)$$

where $I_{|m+\alpha|}$ is the modified Bessel function of the first kind.

Proof. Recall the integral representation

$$I_\nu(z) = \frac{z^\nu}{2^\nu \Gamma(\nu + \frac{1}{2}) \Gamma(\frac{1}{2})} \int_{-1}^1 (1-s^2)^{\nu-\frac{1}{2}} e^{zs} ds, \quad z \in \mathbb{C}, \quad (6.5)$$

see [AS, eq.9.6.18]. Consider now the operator

$$L_m = U h_m U^{-1} \quad \text{in} \quad L^2(\mathbb{R}_+, dr), \quad (6.6)$$

where $U : L^2(\mathbb{R}_+, r dr) \rightarrow L^2(\mathbb{R}_+, dr)$ is the unitary mapping acting as $(Uf)(r) = r^{1/2}f(r)$. Note that L_m is subject to Dirichlet boundary condition at zero and that it coincides with the Friedrichs extension of the differential operator

$$-\frac{d^2}{dr^2} + \frac{\mu^2 - \frac{1}{4}}{r^2}$$

defined on $C_0^\infty(\mathbb{R}_+)$. From [Ko, Sect. 5] we know that

$$W_m L_m W_m^{-1} \varphi(p) = p \varphi(p), \quad \varphi \in W_m(D(L_m)), \quad (6.7)$$

where the mappings $W_m, W_m^{-1} : L^2(\mathbb{R}_+) \rightarrow L^2(\mathbb{R}_+)$ given by

$$(W_m u)(p) = \int_0^\infty u(r) \sqrt{r} J_{|m+\alpha|}(r\sqrt{p}) dr, \quad (W_m^{-1} \varphi)(r) = \frac{1}{2} \int_0^\infty \varphi(p) \sqrt{r} J_{|m+\alpha|}(r\sqrt{p}) dp, \quad (6.8)$$

and defined initially on $C_0^\infty(\mathbb{R}_+)$ extend to unitary operators on $L^2(\mathbb{R}_+)$. By [T, Thm.3.1]

$$e^{-itL_m} g = \lim_{\varepsilon \rightarrow 0+} e^{-(\varepsilon+it)L_m} g, \quad g = Uf. \quad (6.9)$$

In view of (6.7) we thus get

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0+} (e^{-(\varepsilon+it)L_m} g)(r) &= (W_m^{-1} e^{-(\varepsilon+it)p} W_m g)(r) \\ &= \lim_{\varepsilon \rightarrow 0+} \frac{1}{2} \int_0^\infty \sqrt{rr'} \int_0^\infty e^{-(\varepsilon+it)p} J_{|m+\alpha|}(r\sqrt{p}) J_{|m+\alpha|}(r'\sqrt{p}) dp g(r') dr' \\ &= \lim_{\varepsilon \rightarrow 0+} \frac{1}{2(\varepsilon+it)} \int_0^\infty \sqrt{rr'} I_{|m+\alpha|} \left(\frac{rr'}{2(\varepsilon+it)} \right) e^{-\frac{r^2+r'^2}{4(\varepsilon+it)}} g(r') dr'. \end{aligned} \quad (6.10)$$

where we have used [Erd1, Eq.4.14(39)] to evaluate the p -integral. Moreover, from (6.5) it follows that

$$\left| I_{|m+\alpha|} \left(\frac{rr'}{2(\varepsilon+it)} \right) e^{-\frac{r^2+r'^2}{4(\varepsilon+it)}} \right| \leq C \left| \frac{rr'}{t} \right|^{|m+\alpha|} \quad (6.11)$$

for some constant $C > 0$. On the other hand, by assumptions on f we have $(r')^{\frac{1}{2}+|m+\alpha|}g(r') = (r')^{1+|m+\alpha|}f(r') \in L^1(\mathbb{R}_+)$. Hence using the Lebesgue dominated theorem and (6.11) we can interchange the limit with the integral in (6.10) and use (6.9) to get

$$(e^{-itL_m}g)(r) = \frac{1}{2it} \int_0^\infty \sqrt{rr'} I_{|m+\alpha|} \left(\frac{rr'}{2it} \right) e^{-\frac{r^2+r'^2}{4it}} g(r') dr'.$$

Equation (6.4) now follows in view of (6.6) and the identity $g(r') = \sqrt{r'} f(r')$. \square

Now let $u \in L^2(\mathbb{R}^2)$ have compact support. Since the projector Π_m commutes with $h_m \otimes \text{id}$, see section 4, from (4.2) and (6.4) we obtain

$$(e^{-itH_\alpha}u)(r, \theta) = \frac{1}{4\pi it} \sum_{m \in \mathbb{Z}} \int_0^\infty \int_0^{2\pi} I_{|m+\alpha|} \left(\frac{rr'}{2it} \right) e^{im(\theta-\theta')} e^{-\frac{r^2+r'^2}{4it}} u(r', \theta') r' dr' d\theta'. \quad (6.12)$$

Moreover, in view of the integral representation (6.5) the series $\sum_{m \in \mathbb{Z}} I_{|m+\alpha|} \left(\frac{rr'}{2it} \right) e^{im(\theta-\theta')}$ converges, for a fixed r , uniformly in r' on compacts of \mathbb{R}_+ . We can thus interchange the sum and the integral in (6.12) to conclude that (3.1) holds true with

$$e^{-itH_\alpha}(x, y) = \frac{1}{4\pi it} e^{-\frac{r^2+r'^2}{4it}} \sum_{m \in \mathbb{Z}} I_{|m+\alpha|} \left(\frac{rr'}{2it} \right) e^{im(\theta-\theta')}. \quad (6.13)$$

Assume now that $|\alpha| < 1/2$. With the help of (6.5) it is easily seen that

$$\lim_{t \rightarrow \infty} t^{|\alpha|} e^{-\frac{r^2+r'^2}{4it}} \sum_{m \neq 0} I_{|m+\alpha|} \left(\frac{rr'}{2it} \right) e^{im(\theta-\theta')} = 0.$$

Equation (3.2) now follows from the identity

$$\lim_{z \rightarrow 0} z^{-\nu} I_\nu(z) = \frac{1}{2^\nu \Gamma(1+\nu)},$$

see [AS, eq. 9.6.10]. The proof for $|\alpha| = 1/2$ follows the same line. To prove (3.4) we recall that by [FFFP, Corollary 1.7] there exists a constant C_0 such that

$$\sup_{x, y \in \mathbb{R}^2} |e^{-itH_\alpha}(x, y)| \leq \frac{C_0}{t} \quad \forall t > 0. \quad (6.14)$$

Let us define

$$\Omega_1 := \{(x, y) \in \mathbb{R}^4 : |x||y| < t\}, \quad \Omega_2 := \{(x, y) \in \mathbb{R}^4 : |x||y| \geq t\}, \quad z := \frac{|x||y|}{t}.$$

Since $|\alpha| \leq 1/2$, from (6.14), (6.13) and (6.5) we deduce that

$$\begin{aligned} & \sup_{(x, y) \in \mathbb{R}^4} \left(\frac{t}{|x||y|} \right)^{|\alpha|} |e^{-itH_\alpha}(x, y)| \\ &= \max \left\{ \sup_{(x, y) \in \Omega_1} \left(\frac{t}{|x||y|} \right)^{|\alpha|} |e^{-itH_\alpha}(x, y)|, \sup_{(x, y) \in \Omega_2} \left(\frac{t}{|x||y|} \right)^{|\alpha|} |e^{-itH_\alpha}(x, y)| \right\} \\ &\leq t^{-1} \max \left\{ C_1 \sup_{0 < z < 1} \sum_{m \in \mathbb{Z}} \frac{z^{|m+\alpha|-|\alpha|}}{2^{|m+\alpha|} \Gamma(|m+\alpha|+1/2)}, C_0 \right\} = C_2 t^{-1}, \end{aligned}$$

where C_1 and C_2 are positive constants. This implies (3.4) and completes the proof of Theorem 3.1.

Remark 6.3. By using the relation $I_\nu(z) = e^{-\frac{1}{2}i\pi z} J_\nu(iz)$, see [AS, eq.9.6.3], it is easily seen that equation (6.13) agrees with the expression for the Aharonov-Bohm propagator found in [FFFP, Thm. 1.3].

APPENDIX A.

For reader's convenience we recall below the results obtained in [JK, Lemma 10.1] and [JK, Lemma 10.2] regarding a function $F : \mathbb{R}_+ \rightarrow \mathcal{B}$, where \mathcal{B} is an arbitrary Banach space.

Lemma A.1 (Jensen-Kato). *Suppose that $F(\lambda) = 0$ in a neighborhood of zero and that $F'' \in L^1(\mathbb{R}_+; \mathcal{B})$. Then*

$$\int_0^\infty e^{-it\lambda} F(\lambda) d\lambda = o(t^{-2}) \quad \text{as } t \rightarrow \infty \text{ in } \mathcal{B}.$$

Lemma A.2 (Jensen-Kato). *Suppose that $F(0) = 0$, $F(\lambda) = 0$ for λ large enough and that $F'' \in L^1((\delta, \infty); \mathcal{B})$ for any $\delta > 0$. Assume moreover that $F''(\lambda) = o(\lambda^{\beta-2})$ as $\lambda \rightarrow 0$ for some $\beta \in (0, 1)$. Then*

$$\int_0^\infty e^{-it\lambda} F(\lambda) d\lambda = o(t^{-1-\beta}) \quad \text{as } t \rightarrow \infty \text{ in } \mathcal{B}.$$

Remark A.3. [JK] provides more general versions of the above results. Here we limit ourselves to particular situations which suit our purposes.

Acknowledgements

G.G. has been supported by the MIUR-PRIN08 grant for the project "Metodi di viscosit, geometrici e di controllo per modelli diffusivi nonlineari". H.K. has been supported by the MIUR-PRIN08 grant for the project "Trasporto ottimo di massa, disuguaglianze geometriche e funzionali e applicazioni".

REFERENCES

- [AS] M. Abramowitz and I. Stegun, *Handbook of mathematical functions*. National Bureau of Standards, 1964.
- [AT] R. Adami, A. Teta: On the Aharonov-Bohm Hamiltonian. *Lett. Math. Phys.* **43** (1998) 43–53.
- [Erd1] A. Erdelyi: *Tables of integral transforms*, Vol. 2. McGraw-Hill, New York 1954.
- [Erd2] A. Erdelyi: Asymptotic representations of Fourier integrals and the method of stationary phase. *J. Soc. Indust. Appl. Math.* **3** (1955) 17–27.
- [EG] M. B. Erdogan, W. R. Green: A weighted dispersive estimate for Schrödinger operators in dimension two. arXiv: 1202.0050v1. (2012).
- [ES1] M. B. Erdogan, W. Schlag: Dispersive estimates for Schrodinger operators in the presence of a resonance and/or an eigenvalue at zero energy in dimension three: I. *Dyn. Partial Differ. Equ.* **1** (2004) 359–379.
- [ES2] M. B. Erdogan, W. Schlag: Dispersive estimates for Schrodinger operators in the presence of a resonance and/or an eigenvalue at zero energy in dimension three: II. *J. Anal. Math.* **99** (2006) 199–248.
- [FFFP] L. Fanelli, V. Felli, M. A. Fontelos, A. Primo: Time decay of scaling critical electromagnetic Schrödinger flows, arXiv: 1203.1771. (2012).
- [Go] M. Goldberg: Transport in the one-dimensional Schrödinger equation. *Proc. Amer. Math. Soc.* **135** (2007) 3171–3179.
- [GS] M. Goldberg and W. Schlag: Dispersive estimates for Schrödinger operators in dimensions one and three. *Comm. Math. Phys.* **vol. 251** (2004) 157–178.
- [IT] H. T. Ito, H. Tamura: AharonovBohm Effect in Scattering by Point-like Magnetic Fields at Large Separation. *Ann. Henri Poincaré* **2** (2001) 309–359.
- [Je] A. Jensen: Local decay in time of solutions to Schrödingers equation with a dilation-analytic interaction. *Manuscripta Math.* **25** (1978) 61–77.
- [JK] A. Jensen, T. Kato: Spectral properties of Schrödinger operators and time-decay of the wave functions. *Duke Math. J.* **46** (1979) 583–611.
- [JSS] J.L. Journe, A. Soffer, C.D. Sogge: Decay estimates for Schrödinger operators. *Comm. Pure Appl. Math.*, **44** (1991) 573–604.
- [KK] A. I. Komech, E. A. Kopylova: Weighted decay for magnetic Schrödinger equation. arXiv: 1204.1731v1. (2012).
- [Ko] H. Kovařík: Heat kernels of two-dimensional magnetic Schrödinger and Pauli operators. *Calc. Var. Partial Differential Equations* **44** (2012) 351–374.

- [Kr] D. Krejčířík : The improved decay rate for the heat semigroup with local magnetic field in the plane. *Calc. Var. Partial Differential Equations*, in press. DOI 10.1007/s00526-012-0516-1.
- [LW] A. Laptev and T. Weidl: Hardy inequalities for magnetic Dirichlet forms, *Oper. Theory Adv. Appl.* **108** (1999) 299–305.
- [Mu] M. Murata: Asymptotic expansions in time for solutions of Schrödinger-type equations. *J. Funct. Anal.* **49** (1982), 10–56.
- [Ra] J. Rauch: Local decay of scattering solutions to Schrödingers equation. *Comm. Math. Phys.* **61** (1978) 149–168.
- [Sch1] W. Schlag: Dispersive estimates for Schrödinger operators in dimension two. *Comm. Math. Phys.* **257** (2005), 87–117.
- [Sch2] W. Schlag: Dispersive estimates for Schrödinger operators. Mathematical aspects of nonlinear dispersive equations, 255–285, Ann. of Math. Stud., 163, Princeton Univ. Press, Princeton, NJ, 2007.
- [T] G. Teschl: *Mathematical Methods in Quantum Mechanics With Applications to Schrödinger Operators*. American Mathematical Society Providence, Rhode Island, 2009.
- [Wed] R. Weder: $L^p - L^{p'}$ Estimates for the Schrödinger Equation on the Line and Inverse Scattering for the Nonlinear Schrödinger Equation with a Potential. *J. Funct. Anal.* **170** (2000), 37–68.
- [W] T. Weidl: A remark on Hardy type inequalities for critical Schrödinger operators with magnetic fields. *Op. Theory: Adv. and Appl.* **110** (1999) 247–254.
- [Z] V.A. Zorich: *Mathematical Analysis* vol. 2, Springer-Verlag, 2004.

GABRIELE GRILLO, DIPARTIMENTO DI MATEMATICA, POLITECNICO DI MILANO, ITALY
E-mail address: gabriele.grillo@polimi.it

HYNEK KOVAŘÍK, DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI BRESCIA, ITALY
E-mail address: hynek.kovarik@ing.unibs.it